:: Main Page • Future Missions • Feeds • Article Directory • Site Search • About & Quick Guid

« robotic exploration »

Today is Thursday, August 03 2006

Countdown

STEREO launch: 28 days, 4 hours, 1 minutes

SMART-1 lunar

impact: 30 days, 14 hours, 30 minutes

Solar-B launch: 50 days, 6 hours, 21 minutes

Topics

Mars Recon. Orbiter (19)

Stardust (17)

Cassini-Huygens (105)

Mars Exploration Rovers (48)

New Horizons (22)

Deep Impact (10)

Dawn (3)

MESSENGER (2)

Space Observatories (114)

European Space Agency (53)

Space Sci-Tech (72)

Mars (107)

Archive (522)

Upcoming Events

August 31 - STEREO launch.

September 3 -SMART-1 lunar impact. September 7 -



Gigantic cosmic cataclysm in Stephan's Quintet of galaxies Monday, March 06 2006

Recent infrared observations made with Spitzer Space Telescope have revealed the presence of a huge intergalactic shock wave, or "sonic boom" in the middle of Stephan's Quintet, a group of galaxies which is now the scene of a gigantic cosmic cataclysm.



This discovery, made by an international research team including scientists from the Max Planck Institute for Nuclear Physics (MPIK) in Heidelberg, provides a local view of what might have been going on in the early universe, when vast mergers and collisions between galaxies were commonplace.

When astronomers using Spitzer Space Telescope turned their attention to a well-known group of galaxies called Stephan's Quintet, they were, quite simply, shocked at what they saw. There, sweeping through the group, lurks one of the biggest shock waves ever seen.

For decades, astronomers using optical telescopes have known that the galaxies in this group, located about 300 million light years away, have a very distorted distribution of visible light from stars, indicating that the galaxies have experienced encounters in the past, and are now engaged in further collisions. But this, as it turns out, is only part of the drama. Recently, astronomers have become able to measure what, apart from the stars, is present in Stephan's Quintet. By looking in the radio and X-rays they discovered huge quantities of gas - about 100,000 million solar masses, mainly composed of hydrogen and helium - in the space between the galaxies, more than all the gas inside the galaxies themselves.

Now, a team of scientists from Caltech, USA and from the Astrophysics Department of the Max Planck Institute for Nuclear Physics (MPIK) in Heidelberg, Germany, together with other collaborators from the USA and Australia, have turned the Spitzer Space Telescope, equipped with a super-sensitive infrared spectrograph, towards the location of the group. They discovered that one of the galaxies, called NGC7318b, which is falling towards the others at high speed, is generating a giant shock wave in front of it - larger even than the Milky Way - as it ploughs its way through the intergalactic gas. The results of this amazing discovery are to be published on March 10th in a paper in the Astrophysical Journal Letters.

The signature of the shock-wave was given by the detection of strong

Cassini flyby #17 of Titan (Altitude 1,000 km / 600 mi).

September 23 -

Cassini flyby #18 of Titan (Altitude 1,000 km / 600 mi).

September 22 -Solar-B launch.

October 9 - Cassini flyby #19 of Titan (Altitude 1,000 km / 600 mi).

October 19 -THEMIS launch. October 24 -

MESSENGER Venus flyby 1.

October 25 - Cassini flyby #20 of Titan (Altitude 1,000 km / 600 mi).

More...



Older Stories

Thursday 27-Jul

 Mersenius crater wrinkles between Humorum and Procellarum (0)

Wednesday 26-Jul

- Census conducted of nearby hidden black holes (0)
- Old pulsars still have new tricks

radiation from molecular hydrogen. When hydrogen molecules are "excited" by the mechanical energy produced in the collision and transported by shock waves, they emit a distinctive type of radiation that can be detected in the infrared, and it was this radiation that was picked up by Spitzer. "The strength of the emission and the fact that it shows the gas to be highly disturbed was a huge surprise to us, said team leader Dr. Phil Appleton. "We expected to see the spectral signature of dust grains - but instead we saw an almost pure laboratory-like spectrum of hydrogen molecules and almost nothing else. It was quite unlike anything we had seen before in a galaxy system."



Image Credit: NASA/JPL-Caltech/Max-Planck Institute/P. Appleton (SSC/Caltech)

This false-color composite image of the Stephan's Quintet galaxy cluster clearly shows one of the largest shock waves ever seen (green arc), produced by one galaxy falling toward another at over a million miles per hour. It is made up of data from Spitzer Space Telescope and a ground-based telescope in Spain.

Four of the five galaxies in this image are involved in a violent collision, which has already stripped most of the hydrogen gas from the interiors of the galaxies. The centers of the galaxies appear as bright yellow-pink knots inside a blue haze of stars, and the galaxy producing all the turmoil,

to teach us (0)

 Mars Global Surveyor images - July 20-26, 2006 (0)

Tuesday 25-Jul

- Lakes on Titan (0)
- Titan's pebbles 'seen' by Huygens radio (0)
- Lomonosov a large crater filled by lava (0)

Monday 24-Jul

- SMART-1 birthday postcard of Apollo 11 landing site (0)
- Planet-forming disks might put the brakes on stars (0)

Saturday 22-Jul

 Rock abrasion tool exhibits the deep red pigment of Mars (0)

User Functions

Username:

Password:

Login

Don't have an account yet? Sign up as a **New User** NGC7318b, is the left of two small bright regions in the middle right of the image. One galaxy, the large spiral at the bottom left of the image, is a foreground object and is not associated with the cluster.

The titanic shock wave, larger than our own Milky Way galaxy, was detected by the ground-based telescope using visible-light wavelengths. It consists of hot hydrogen gas. As NGC7318b collides with gas spread throughout the cluster, atoms of hydrogen are heated in the shock wave, producing the green glow.

Spitzer pointed its infrared spectrograph at the peak of this shock wave (middle of green glow) to learn more about its inner workings. This instrument breaks light apart into its basic components. Data from the instrument are referred to as spectra and are displayed as curving lines that indicate the amount of light coming at each specific wavelength.

The Spitzer spectrum showed a strong infrared signature for incredibly turbulent gas made up of hydrogen molecules. This gas is caused when atoms of hydrogen rapidly pair-up to form molecules in the wake of the shock wave. Molecular hydrogen, unlike atomic hydrogen, gives off most of its energy through vibrations that emit in the infrared.

This highly disturbed gas is the most turbulent molecular hydrogen ever seen. Astronomers were surprised not only by the turbulence of the gas, but by the incredible strength of the emission. The reason the molecular hydrogen emission is so powerful is not yet completely understood.

Stephan's Quintet is located 300 million light-years away in the Pegasus constellation.

This image is composed of three data sets: near-infrared light (blue) and visible light called H-alpha (green) from the Calar Alto Observatory in Spain, operated by the Max Planck Institute in Germany; and 8-micron infrared light (red) from Spitzer's infrared array camera.

Spectrometers have the ability to break light down into its component wavelengths, where the chemical signatures of the material that produced it can be seen as spectral lines. The width of these lines allows astronomers to determine the velocity of the gas, with wider lines indicating gas at a higher velocity. The hydrogen line seen by Spitzer in Stephan's Quintet is the widest ever observed for hot hydrogen molecules, corresponding to gas motions of 870 kilometres per second, equivalent to a Mach number of 100 or more. "To better understand this situation", says Dr. Richard Tuffs, a team member from MPIK, "one can think of the shock waves created by supersonic aircraft in the Earth atmosphere. There, water droplets can condense behind the shock in humid conditions, while in Stephan's Quintet hydrogen molecules could form out of a turbulent and cooling intergalactic medium. Of course, all this is happening on an enormous scale in Stephan's Quintet."

The Spitzer observations provide a new diagnostic for studying conditions in merging and colliding galaxies, which were much more prevalent in the early universe. "Observing a nearby densely populated galaxy group, immersed in a thick gas cloud, gives us a local view of what might have been going on in the early universe about 10 billion years ago, soon after the first galaxies formed, when the intergalactic medium and the galaxy density were much greater than today. In this respect these observations are a bit like stepping into a time machine", said Dr. Cristina Popescu, another team member from MPIK. The new results may indicate that some of the emission from the most luminous infrared galaxies seen in the very distant Universe may actually be created not by stars, but by vast shocks in the gas between the colliding galaxies.

Though far in the future, it is likely that in about two billion years from now, we will collide with the slightly larger neighbouring Andromeda Galaxy, creating shocks of our own.

NASA's Jet Propulsion Laboratory, Pasadena, California, manages the Spitzer Space Telescope mission for the agency's Science Mission Directorate. Science operations are conducted at the Spitzer Science Center at Caltech. JPL is a division of Caltech.

Max Planck Society News Release

1000%+ Stock AlertsLes infos temps réelThe Stockster has twoMes quotidiens sur mon
Gigantic cosmic cataclysm in Stephan's Quintet of galaxies 0 Comments Create New Account
Oldest First Threaded Refresh Reply
The following comments are owned by whomever posted them. This site is not responsible for what they say.
Privacy Policy
Created this page in 0.28 seconds