To see related records in other databases, click on the icons below (if any are shown) or on the links highlighted within the body of this record (if any are present).

New model helps to calculate energy output of stars

[Date: 2008-05-20]

Category: Miscellaneous

Data Source: Max Planck Institute for Nuclear Physics; The Astrophysical Journal

Provider: Driver, Simon P. et al. (2008) The energy output of the Universe from 0.1 to 1000 micrometres. The Astrophysical Journal, 678:L101-L104, 10 May 2008.

Document of reference:

Subject index: Coordination, Cooperation, Scientific Research, Space & Satellite research

Programme Acronym: MSD-FC MS-UK-C



The observations are based on a new model developed by Dr Cristina Popescu of the University of Central Lancashire in the UK and Dr Richard Tuffs of the Max Planck Institute for Nuclear Physics in Heidelberg, Germany. The model describes how dust particles are distributed in relation to stars and disks of galaxies and, by calculating the dimming effect of the dust, makes it possible to predict the difference between the total light emitted by the stars and the total light inferred from observations.

An average star at a random position in the universe generates about 40 quadrillion kilowatthours in one year. This amounts to 300 times humankind's current energy consumption needs.

Similar to smoke in the Earth's atmosphere, dust particles absorb large parts of the energy emitted by stars. Of course, the energy is not lost, but it heats the dust particles, which emit heat radiation as a result. This heat radiation can be detected in the infrared part of the electromagnetic spectrum.

The amount of energy absorbed by the particles is determined by their position in relation to the stars, which is why the model developed by Dr Popescu and Dr Tuffs has greatly helped the understanding of the phenomenon. In previous calculations, the infrared emissions of dust sometimes seemed to surpass estimates of total starlight emitted. 'You can't get more energy out than you put in - we knew that something wasn't quite right there,' team leader Dr Simon Driver of the University of St Andrews, UK, explains. A paradox that the new model has resolved.

'The fact that the observations so closely matched the predictions make us confident that we really now understand the radiation of energy from galaxies and thereby from the universe as a whole over a wide range of wavelengths,' says Dr Popescu.

'The results very clearly show that interstellar dust particles have a very pronounced effect on the observed starlight from galaxies in the local universe,' adds Dr Tuffs.

The researchers hope to soon get an even more detailed picture of individual galaxies with the help of two new facilities due to be up and running in the coming months, namely the VISTA telescope in Chile and 'Herschel', the European Space Agency's (ESA's) infrared space observatory. 'VISTA will allow us to see further through the dust, whereas Herschel will directly detect the emission from the dust,' says Dr Jochen Liske of the European Southern Observatory.